arXiv:1206.1937v1 [math.GT] 9 Jun 2012

FINDING A SYSTEM OF ESSENTIAL 2-SUBORBIFOLDS

YOSHIHIRO TAKEUCHI AND MISAKO YOKOYAMA

Abstract. We make an analogy of Culler-Morgan-Shalen theory. Our main goal is to show that there exists a non-empty system of essential 2-suborbifolds respecting a given splitting of the orbifold fundamental group.

MSC:primary 57M50; secondary 57M60

1. Introduction

For a 3-dimensional manifold N, the essential (i.e., incompressible and not boundary parallel) 2-suborbifolds are corresponding to the decompositions of the fundamental group of N. If N has an essential and separating 2-subsphere, $\pi_1(N)$ has the free product decomposition which respects its geometric decomposition, and conversely, if $\pi_1(N)$ has a free product decomposition, N has an essential and separating 2-subsphere which realizes its algebraic decomposition. If N has an essential 2-submanifold S which is not a 2-sphere but separating, the above decomposition of $\pi_1(N)$ turns to be an amalgamated free product decomposition, and if S is non-separating, it does to be an HNN extension decomposition. Moreover, [C-S] proved the theorem that if N acts on a simplicial tree nontrivially, N has a non-empty system of essential 2-submanifolds which respects that action.

A similar approach should be considered for 3-orbifolds. If a 3-obifold M has an essential 2-suborbifold, it is clear that the orbifold fundamnetal group $\pi_1(M)$ has an amalgamated free product decomposition or an HNN extension decomposition. In [T-Y 2] (respectively, [T-Y 3]) we found an essential non-spherical (respectively, spherical) 2-suborbifold realizing a given algebraic decomposition of the orbifold fundamental group of M.

In the present paper we show the following:

Theorem 1.1. *Let M be a good, compact, connected, orientable 3-orbifold without non-separating spherical 2-orbifolds. We assume that the fundamental group of each prime component of M is infinite. Suppose that $\pi_1(M)$ has a nontrivial finite splitting. Then there exists a non-empty system of essential 2-suborbifolds $S_1, \ldots, S_n \subset M$ such that for each component Q of $M - \bigcup_{i=1}^{n} S_i$, $\pi_1(Q)$ is contained in a vertex group.*

Boileau, Maillot and Porti showed a related result in [BMP, Proposition 7.16], where they treat with the fundamental group of the complement of the set of singular points of a 3-orbifold.

We summarize the contents of the present paper. In Section 2, 3 and 4, we review on the actions on a tree, 3-orbifolds, and OISIBO's (orbifold identified spaces

---

Key words and phrases. orbifold, orbifold fundamental group, Culler-Morgan-Shalen theory.

identified along ballic orbifolds) respectively. In Section 5, we prepare an orbifold composition, which is used in Main Theorem as the target space of a b-continuous map. In Section 6, we prove Main Theorem.

2. Preliminaries on the actions on a tree

Throughout this present paper any orbifold is assumed to be good, that is, it is covered by a manifold, and assumed to be compact, connected and orientable unless otherwise stated.

In [Se], some fixed point theorems about group actions on trees are proved. Here we use the following restricted forms of them.

Let T be a simplicial tree, i.e., a connected and simply connected 1-complex, and G a group simplicially acting on T.

For $g \in G$, g is called to have an *edge inversion* if there exists an edge E such that $g(E) = E$ and $g|E$ is orientation reversing.

The action is called *trivial* if a vertex of T is fixed by Γ.

Proposition 2.1. *Let g be an element of G with finite order. If g acts on T without edge inversions, then there exists a vertex p of T such that $g(p) = p$.*

Proposition 2.2. *Let p_1, $p_2 \in T$ be fixed points of $g \in G$ and ℓ the unique simple path from p_1 to p_2. Then any vertex and edge on ℓ are fixed by g.*

Let $n \geq 1$ be an integer. Put

$$G_n = \langle\, a_1, \cdots, a_n \,|\, a_1^{\alpha_1} = \cdots = a_n^{\alpha_n} = (a_i a_j)^{\beta_{i,j}} = 1,\ 1 \leq i < j \leq n \,\rangle$$

where $\alpha_i, \beta_{i,j} \geq 2$ are integers.

Lemma 2.3. *If G_n acts on T without edge inversions, then T has a fixed vertex of G_n action.*

3. Preliminaries on orbifolds

Definition 3.1. Let $M = (\tilde{M}, p, |M|)$, $N = (\tilde{N}, q, |N|)$ be orbifolds. A *continuous map* $f : M \to N$ is a pair $(|f|, \tilde{f})$ of continuous maps $|f| : |M| \to |N|$ and $\tilde{f} : \tilde{M} \to \tilde{N}$ which satisfies the following:

$$\begin{tikzcd} \tilde{M} \arrow[r, "\tilde{f}"] \arrow[d, "p"'] & \tilde{N} \arrow[d, "q"] \\ {|M|} \arrow[r, "{|f|}"'] & {|N|} \end{tikzcd} \tag{3.1}$$

- **(i)** $|f| \circ p = q \circ \tilde{f}$,
- **(ii)** For each $\sigma \in \mathrm{Aut}(\tilde{M}, p)$ there exists an element $\tau \in \mathrm{Aut}(\tilde{N}, q)$ such that $\tilde{f} \circ \sigma = \tau \circ \tilde{f}$.

A continuous map $f : M \to N$ is *b-continuous* if there exists a point $x \in |M| - \Sigma M$ such that $|f|(x) \in |N| - \Sigma N$. It was called an orbi-map in [Ta1, T-Y 1], etc. A b-continuous map induces a homomorphism between the fundamental groups and local fundamental groups of orbifolds, see [T-Y 4, Lemma 3.13] and [Ta1], where the points x and $|f|(x)$ in the above could be base points of the fundamental groups of M and N, respectively. The notion of (b-)continuous maps between orbifolds is naturally generalized for those between OISIBO's and orbifold compositions in Sections 4 and 5.

A b-continuous map $f : M \to N$ is an *embedding* if $f(M)$ is a suborbifold of N and $f : M \to f(M)$ is an isomorphism of orbifolds.

For other terminologies, see [Ta1].

4. Preliminaries on OISIBO's

Definition 4.1. Let I, J be countable sets, X_i $(i \in I)$ n-orbifolds, and B_j $(j \in J)$ ballic n-orbifolds. Let $f_j^\varepsilon : B_j \to X_{i(j,\varepsilon)}$ be embeddings (as orbifolds) such that $f_j^\varepsilon(B_j) \subset \text{Int } X_{i(j,\varepsilon)}$ and $f_j^\varepsilon(B_j)$ are mutually disjoint, where $j \in J$, $i(j,\varepsilon) \in I$, $\varepsilon = 0, 1$. Then we call $X = (X_i, B_j, f_j^\varepsilon)_{i \in I, j \in J, \varepsilon = 0,1}$ an *n-orbifold identified space identified along ballic orbifolds* (n-OISIBO). The maps $f_j^0 \circ (f_j^1)^{-1}$ and their inverses are called the *identifying maps* of X. Each X_i, B_j are called a *particle of* X, and an *identifying ballic orbifold*, respectively. We define the equivalence relation $\sim$ in $\coprod_{i \in I, j \in J}(|X_i| \cup |B_j|)$ to be generated by

$$y \sim |f|_j^\varepsilon(y), \quad \varepsilon = 0, 1, \quad y \in |B_j|, \quad j \in J. \tag{4.1}$$

We call the identified space $\coprod_{i \in I, j \in J}(|X_i| \cup |B_j|)/\sim$ the *underlying space of* X, denoted by $|X|$, and call the identified space $\{(\cup_{i \in I}\Sigma X_i) \cup (\cup_{j \in J}\Sigma B_j)\}/\sim$ the *singular set of* X, denoted by ΣX.

Definition 4.2. Let $X = (X_i, B_j, f_j^\varepsilon)_{i \in I, j \in J, \varepsilon = 0,1}$ and $X' = (X'_k, B'_\ell, g_\ell^\varepsilon)_{k \in K, \ell \in L, \varepsilon = 0,1}$ be OISIBO's. We say that X and X' are *isomorphic* if there exists a set of maps $\{\varphi_i, \psi_j\}_{i \in I, j \in J}$ and bijections $\eta : I \to K$, $\xi : J \to L$ such that the following (i) and (ii) hold:

- **(i)** For each $i \in I$, φ_i is an isomorphism (of orbifolds) from X_i to $X'_{\eta(i)}$, and for each $j \in J$, ψ_j is an isomorphism (of orbifolds) from B_j to $B'_{\xi(j)}$.
- **(ii)** For each $j \in J$, and $\varepsilon = 0, 1$, $\varphi_{i(j,\varepsilon)} \circ f_j^\varepsilon = g_{\xi(j)}^\varepsilon \circ \psi_j$.

The system $h = (\{\varphi_i, \psi_j\}_{i \in I, j \in J}, \eta, \xi)$ is called an *isomorphism from X to X'*.

Definition 4.3. Let $X = (X_k, B_\ell, f_\ell^\varepsilon)_{k \in K, \ell \in L, \varepsilon = 0,1}$ and $X' = (X'_i, B'_j, {f'}_j^\varepsilon)_{i \in I, j \in J, \varepsilon = 0,1}$ be OISIBO's. We say that X' is a *covering of* X if there exists a set of maps $\{\varphi_i, \psi_j\}_{i \in I, j \in J}$ and surjections $\eta : I \to K$, $\xi : J \to L$ such that the following (i) and (ii) hold:

- **(i)** Each φ_i is a covering map (of orbifolds) from X'_i to $X_{\eta(i)}$, where $\eta(i) \in K$, and each ψ_j is a covering map (of orbifolds) from B'_j to $B_{\xi(j)}$, where $\xi(j) \in L$.
- **(ii)** For each $j \in J$ and $\varepsilon = 0, 1$, $\varphi_{i(j,\varepsilon)} \circ {f'}_j^\varepsilon = f_{\xi(j)}^\varepsilon \circ \psi_j$.

Note that the continuous map $|p| : |X'| \to |X|$ naturally induced by $\{\varphi_i, \psi_j\}_{i \in I, j \in J}$ is surjective, and induces the usual covering map from $|X'| - |p|^{-1}(\Sigma X)$ to $|X| - \Sigma X$. We call the system $p = (|p|, \{\varphi_i, \psi_j\}_{i \in I, j \in J})$ a *covering map from X' to X*.

Definition 4.4. Let $\tilde{X}$, X be OISIBO's, and $p : \tilde{X} \to X$ a covering. We call p a *universal covering* if for any covering $p' : X' \to X$, there exists a covering $q : \tilde{X} \to X'$ such that $p = p' \circ q$. As the usual covering theory, for any OISIBO X, there exists a unique universal covering $p : \tilde{X} \to X$.

Definition 4.5. Let X', X be OISIBO's, and $p : X' \to X$ a covering. We define the *deck transformation group* $\mathrm{Aut}(X', p)$ *of* p by

$$\mathrm{Aut}(X', p) = \{h : X' \to X' \mid h \text{ is an isomorphism such that } p \circ h = p\}. \tag{4.2}$$

We sometimes denote an OISIBO X by $(\tilde{X}, p, |X|)$, where $p : \tilde{X} \to X$ is the universal covering and $|X|$ is the underlying space of X. Any orbifold is considered as a special case of an OISIBO.

Definition 4.6. Let $X = (\tilde{X}, p, |X|)$, $Y = (\tilde{Y}, q, |Y|)$ be OISIBO's. A *continuous map* $f : X \to Y$ is a pair $(|f|, \tilde{f})$ of continuous maps $|f| : |X| \to |Y|$ and $\tilde{f} : \tilde{X} \to \tilde{Y}$ which satisfies the same property as (i) and (ii) in Definition 3.1. A continuous map $f : X \to Y$ is *b-continuous* if there exists a point $x \in |X| - \Sigma X$ such that $|f|(x) \in |Y| - \Sigma Y$.

We define a *homotopy* of OISIBO's by using of continuous maps of OISIBO's as the usual homotopy. If the continuous maps at 0 *and* 1 levels of the homotopy are b-continuous, this homotopy is called a b-*homotopy.* See [T-Y 4].

We define a *path* in an OISIBO X by using of a b-continuous map $\alpha = (|\alpha|, \tilde{\alpha}) : [0, 1] \to X$ with $|\alpha|(0) \in |X| - \Sigma X$. If a path α in X satisfies that $|\alpha|(0) = |\alpha|(1)$, it is called a *loop* in X.

By using of loops in an OISIBO X, we define the fundamental group of X as the usual theory. A b-continuous map $f : X \to Y$ between OISIBO's X and Y induces a homomorphism between the fundamental groups and local fundamental groups of X and Y, where the points $x \in |X| - \Sigma X$ and $|f|(x) \in |Y| - \Sigma Y$ in the definition of b-continuous map are the base points of the fundamental groups of X and Y, respectively.

As usual covering theory, various similar results holds such as the following:

Proposition 4.7. *Let X be an OISIBO and let x, y be any two points of $|X| - \Sigma X$. Then the fundamental groups $\pi_1(X, x)$ and $\pi_1(X, y)$ are isomorphic.*

We often denote $\pi_1(X, x_0)$ by $\pi_1(X)$ dropping a base point if not necessary.

Proposition 4.8. *Let X be an OISIBO and $p : \tilde{X} \to X$ the universal covering of X. Then the fundamental group $\pi_1(X)$ is isomorphic to the deck transformation group* $\mathrm{Aut}(\tilde{X}, p)$.

Proposition 4.9. *Let X be an OISIBO. For each subgroup H of $\pi_1(X)$, there exists a covering $p : \tilde{X} \to X$ such that the OISIBO $\tilde{X}$ has a fundamental group which is isomorphic to H.*

5. Orbifold compositions

Definition 5.1. Let I, J be countable sets, X_i $(i \in I)$ and Y_j $(j \in J)$ be n-OISIBO's. Let $f_j^\varepsilon : Y_j \times \varepsilon \to X_{i(j,\varepsilon)}$ be b-continuous-maps, $f_j^\varepsilon = (|f_j^\varepsilon|, \tilde{f}_j^\varepsilon)$, such that $(f_j^\varepsilon)_*$ are monic, where $j \in J$, $i(j, \varepsilon) \in I$, $\varepsilon = 0, 1$. Then we call $X = (X_i, Y_j \times [0, 1], f_j^\varepsilon)_{i \in I, j \in J, \varepsilon = 0,1}$ an *n-dimensional orbifold composition* (*of type III*). The maps f_j^ε are called the *attaching maps* of X, which may have intersections and self-intersections. Each X_i, $Y_j \times [0, 1]$ is called a *component of* X. The equivalence relation $\sim$ in $\coprod_{i \in I, j \in J}(|X_i| \cup (|Y_j| \times [0, 1]))$ is defined to be generated by

$$(y, \varepsilon) \sim |f_j^\varepsilon|(y, \varepsilon), \quad \varepsilon = 0, 1, \quad y \in |Y_j|, \quad j \in J. \tag{5.1}$$

We call the identified space $\coprod_{i\in I, j\in J}\bigl(|X_i|\cup(|Y_j|\times[0,1])\bigr)/\sim$ the *underlying space of* X, denoted by $|X|$, and call the identified space $\{(\cup_{i\in I}\Sigma X_i)\cup(\cup_{j\in J}\Sigma(Y_j\times[0,1]))\}/\sim$ the *singular set of* X, denoted by ΣX.

In the definition of an orbifold composition, each $(f_j^\varepsilon)_*$ is monic, so that we can obtain the unique lift of any path ℓ in X such that $\ell[0,1)\cap\Sigma X=\emptyset$.

We define a covering of an orbifold composition as similar as that of an OISIBO. We sometimes denote an orbifold composition X by $(\tilde{X}, p, |X|)$, where $p:\tilde{X}\to X$ is the universal covering and $|X|$ is the underlying space of X. Any orbifold and any OISIBO are considered as special cases of an orbifold composition.

Definition 5.2. Let $X=(\tilde{X},p,|X|)$, $Y=(\tilde{Y},q,|Y|)$ be orbifold compositions. A *continuous map* $f:X\to Y$ is a pair $(|f|,\tilde{f})$ of continuous maps $|f|:|X|\to|Y|$ and $\tilde{f}:\tilde{X}\to\tilde{Y}$ which satisfies the same property as (i) and (ii) in Definition 3.1. A continuous map $f:X\to Y$ is *b-continuous* if there exists a point $x\in|X|-\Sigma X$ such that $|f|(x)\in|Y|-\Sigma Y$.

By using of loops in an orbifold composition X, we define the fundamental group of X as the usual theory. A b-continuous map $f:X\to Y$ between orbifold compositions X and Y induces a homomorphism between the fundamental groups and local fundamental groups of X and Y, where the points $x\in|X|-\Sigma X$ and $|f|(x)\in|Y|-\Sigma Y$ in the definition of b-continuous map are the base points of the fundamental groups of X and Y, respectively.

Definition 5.3. Let X be an orbifold composition. Define $O_i(X)$, $i=1,2,3$ as follows:

$$O_1(X)=\{f:S^1\to X\text{, a b-continuous map}\mid[f]\text{ is of finite order }(\neq 1)\text{ in }\pi_1(X)\},$$
$$O_2(X)=\{f:S\to X\text{, a b-continuous map}\mid S\text{ is a spherical 2-orbifold}\},$$
$$O_3(X)=\{f:\mathcal{D}B\to X\text{, a b-continuous map}\mid\mathcal{D}B\text{ is the double of a ballic 3-orbifold }B\}.$$

A b-continuous map $f:S^1\to X\in O_1(X)$ is *trivial* if there exists a b-continuous map g from a discal 2-orbifold D to X such that $g|\partial D=f$ and the index of D equals to the order of $[f]$. $O_1(X)$ is *trivial* if every element of $O_1(X)$ is trivial. We call $f:S\to X\in O_2(X)$ *trivial* if there exists a b-continuous map $g:c*S\to X$ such that $g|S=f$, where $c*S$ is the cone on S. $O_2(X)$ is *trivial* if every element of $O_2(X)$ is trivial. We define the triviality of $O_3(X)$ similarly.

Note that if $O_i(X)$ is trivial, then any covering $\tilde{X}$ of X inherits the triviality.

Remark 5.4. Let X be an orbifold composition, and $\tilde{X}$ the universal cover of X. If $O_2(X)$ is trivial, then $\pi_2(\tilde{X})=0$.

Proposition 5.5. *Let M be a compact 3-orbifold, and X an orbifold composition. If $O_1(X)$ and $O_2(X)$ are trivial, then for any homomorphism $\varphi:\pi_1(M)\to\pi_1(X)$, there exists a b-continuous map $f:M\to X$ such that $f_*=\varphi$.*

Proof. Let $M_0=|M|-$ Int $U(\Sigma M)$, where $U(\Sigma M)$ is the small regular neighborhood of ΣM. We can construct a b-continuous map from the 1-skelton of M_0 to X associated with φ. Since $O_1(X)$ and $O_2(X)$ are trivial, this b-continuous map is extendable to M_0 and furthermore to M, that is, we have obtained the desired b-continuous map. □

Proposition 5.6. *Let X be an orbifold composition, and $f : S^1 \to X$ a b-continuous map. If* $\mathrm{Fix}([f]_A) \neq \phi$*, then f is extendable to a b-continuous map from a discal 2-orbifold D to X where $D = D^2(n)$, and n is the order of $[f]_A$.*

Proof. Let $q : D^2 \to D$ be the universal covering. Choose a point $x \in \mathrm{Fix}([f]_A)$. We can construct the structure map of the desired b-continuous map by mapping the cone point of D^2 to x and performing the skeletonwise and equivariant extension. □

Let S be a spherical 2-orbifold and let $q : (\tilde{S}, \tilde{x}_0) \to (S, x_0)$, $x_0 \notin \Sigma S$, be the universal covering. Let τ be an element of $\pi_1(S, x_0)$, $\tilde{x}_\tau$ one of the two points of $\mathrm{Fix}(\tau_A)$, and $x_\tau = q(\tilde{x}_\tau)$. By the symbol $\mu(x_\tau)$, we shall mean the local normal loop around x_τ. Let ℓ be a path in $|S| - \Sigma S$ from $\mu(x_\tau)(0)$ to x_0 such that $\tau = ([\ell^{-1} \cdot \mu(x_\tau) \cdot \ell])^k$, $k \in \mathbb{Z}$

Proposition 5.7. *Let S be a spherical 2-orbifold, X an orbifold composition, and $f : S \to X$ a b-continuous map. Suppose that there exists a point $\tilde{d} \in$* $\mathrm{Fix}(f_*\pi_1(S))_A$*, and for any $\tau \in \pi_1(S, x_0)$ there exists an interval ℓ_σ including $\tilde{d}$ and $\tilde{f}(\tilde{x}_\tau)$ which is fixed by σ_A, where $\sigma = f_*(\tau)$. If π_2 of the universal cover $\tilde{X}$ of X is 0, then f is extendable to a b-continuous map from the cone on S to X.*

Proof. The proof is similar to that of [T-Y 3, Proposition 5.6]. □

Proposition 5.8. *Let $\mathcal{D}B$ be the double of a ballic 3-orbifold B, X an orbifold composition, and $f : \mathcal{D}B \to X$ a b-continuous map. Suppose that* $\mathrm{Fix}(f_*\pi_1(\partial B))_A$ *is connected, and for $\tau \in \pi_1(\partial B, x_0)$, $\pi_1(\mathrm{Fix}(f_*((\tau))_A)) = 1$ and there exists an interval ℓ_σ including $\tilde{d}$ and $\tilde{f}(\tilde{x}_\tau)$ which is fixed by σ_A, where $\sigma = f_*(\tau)$. If π_2 and π_3 of the universal cover $\tilde{X}$ of X is 0, then f is extendable to a b-continuous map from the cone on $\mathcal{D}B$ to X.*

Proof. The proof is similar to that of [T-Y 3, Proposition 5.7]. □

Proposition 5.9. [T-Y 3, Proposition 5.8] *Let X be a 3-OISIBO whose particles are irreducible. Let $p : \tilde{X} \to X$ be the universal covering and $\sigma \in$* $\mathrm{Aut}(\tilde{X}, p)$ *be any nontrivial element of finite order. Suppose that each particle of $\tilde{X}$ is non-compact. Then the following holds:*

- **(i)** $\mathrm{Fix}(\sigma) \neq \phi$ *and is homeomorphic to a tree.*
- **(ii)** *$O_1(X)$ is trivial.*

Since we assume that any orbifold is orientable, the restriction of σ to each particle is orientation preserving, and each identifying ballic orbifold is orientable.

Proposition 5.10. [T-Y 3, Proposition 5.9] *Let X be a 3-OISIBO, each particle of which is irreducible, and $p : \tilde{X} \to X$ the universal covering. Let G be any subgroup of* $\mathrm{Aut}(\tilde{X}, p)$*, which is isomorphic to the orbifold fundamental group of a spherical 2-orbifold S. Suppose that each particle of $\tilde{X}$ is non-compact. Then the following holds:*

- **(i)** $\mathrm{Fix}(G)$ *is either a point or a tree.*
- **(ii)** $\pi_2(\tilde{X}) = \pi_3(\tilde{X}) = 0$.
- **(iii)** *$O_i(X)$'s are trivial, $i = 1, 2, 3$.*

Proposition 5.11. *Let* $X = (\tilde{X}, p, |X|)$, $Y = (\tilde{Y}, q, |Y|)$ *be orbifold compositions, and* $f = (|f|, \tilde{f}) : X \to Y$ *a b-continuous map. Then for* $[\alpha] \in \pi_1(X, x)$,

$$\tilde{f} \circ [\alpha]_A = (f_*([\alpha]))_A \circ \tilde{f}. \tag{5.2}$$

Proof. Let $\tilde{x} \in p^{-1}(x)$ be the base point of $\tilde{X}$. Note that $[\alpha]_A$ is characterized as the element of $\mathrm{Aut}(\tilde{X}, p)$ which transforms $\tilde{\alpha}(0) = \tilde{x}$ to $\tilde{\alpha}(1)$, and $(f_*([\alpha]))_A = [f \circ \alpha]_A$ is characterized as the element of $\mathrm{Aut}(\tilde{Y}, q)$ which transforms $\tilde{f}(\tilde{\alpha}(0))$ to $\tilde{f}(\tilde{\alpha}(1))$. By the definition of b-continuous map, there exists an element $\tau \in \mathrm{Aut}(\tilde{Y}, q)$ such that $\tilde{f} \circ [\alpha]_A = \tau \circ \tilde{f}$. On the other hand, $\tau(\tilde{f}(\tilde{\alpha}(0))) = (\tau \circ \tilde{f})(\tilde{\alpha}(0)) = (\tilde{f} \circ [\alpha]_A)(\tilde{\alpha}(0)) = \tilde{f}(\tilde{\alpha}(1))$. Hence $\tau = (f_*([\alpha]))_A$. □

Proposition 5.12. *Let* $X = (X_i, Y_j \times [0, 1], f^j_\varepsilon)_{i \in I, j \in J, \varepsilon = 0,1}$ *be an orbifold composition, where each particle of each* X_i *and* Y_j *is an orientable and irreducible 3-orbifold whose universal covering is noncompact. Then* $O_i(X)$*'s are trivial,* $i = 1, 2, 3$.

Proof. Let $p : \tilde{X} \to X$ be the universal covering. From the uniqueness of the universal covering, we may assume that $\tilde{X}$ is the orbifold composition constructed in a similar method described in [T-Y 2].

Claim: Let G be any subgroup of $\mathrm{Aut}(\tilde{X}, p)$, which is isomorphic to the fundamental group of a spherical 2-orbifold. Consider the associated 1-complex $\mathcal{C}(\tilde{X})$ of $\tilde{X}$. Then, there exists a vertex OISIBO $\tilde{Z}$ of $\tilde{X}$ with respect to $\mathcal{C}(\tilde{X})$ such that $G(\tilde{Z}) = \tilde{Z}$.

Indeed, G is finite and acts on the tree $\mathcal{C}(\tilde{X})$ without edge inversions. By Lemma 2.3 we have the claim.

Take any element $f \in O_1(X)$. By the claim, there exists an OISIBO $\tilde{Z}$ of $\tilde{X}$ such that $[f]_A(\tilde{Z}) = \tilde{Z}$. Then by Proposition 5.9, $\mathrm{Fix}([f]_A) \neq \emptyset$ in $\tilde{Z}$. Thus $\mathrm{Fix}([f]_A) \neq \emptyset$ in $\tilde{X}$. By using of Proposition 5.6, f is extendable to a b-continuous map from $D^2(n)$ to X.

Take any element $f \in O_2(X)$, $f : S \to X$. Let $q : \tilde{S} \to S$ be the universal covering and $\tilde{f} : \tilde{S} \to \tilde{X}$ the structure map of f. Let $B = c * S$ be the cone on S and c the cone point of B. Let $\bar{q} : \tilde{B} = \tilde{c} * \tilde{S} \to B$ be the universal covering, $\tilde{c} = \bar{q}^{-1}(c)$ and $\bar{q}(t\tilde{x} + (1-t)\tilde{c}) = tq(\tilde{x}) + (1-t)c$, $\tilde{x} \in \tilde{S}$. Note that $f_*\pi_1(S)$ is isomorphic to a spherical 2-orbifold group. By the claim, there exists a vertex OISIBO $\tilde{Z}$ of $\tilde{X}$ such that $(f_*\pi_1(S))(\tilde{Z}) = \tilde{Z}$. By Proposition 5.10, $\mathrm{Fix}(f_*\pi_1(S))_A \neq \emptyset$. Thus there exists a point $\tilde{d} \in \tilde{Z}$ such that $(f_*\pi_1(S))_A \tilde{d} = \tilde{d}$.

Choose any $\tau \in \pi_1(S)$. We put $\sigma = f_*(\tau)$. Since $\sigma_A \in (f_*\pi_1(S))_A$, $\sigma_A(\tilde{d}) = \tilde{d}$. Moreover, by the fact $\tau_A(\tilde{x}_\tau) = \tilde{x}_\tau$ and Proposition 5.11, $\sigma_A(\tilde{f}(\tilde{x}_\tau)) = f_*(\tau)_A(\tilde{f}(\tilde{x}_\tau)) = \tilde{f} \circ \tau_A(\tilde{x}_\tau) = \tilde{f}(\tilde{x}_\tau)$.

Let $\tilde{Z}_1$ be the OISIBO in which $\tilde{f}(\tilde{x}_\tau)$ is included and let $\tilde{Z}_k$ be the OISIBO in which $\tilde{d}$ is included. Since σ_A acts on the tree $\mathcal{C}(\tilde{X})$, $\tilde{Z}_1$ and $\tilde{Z}_k$ are invariant by σ_A. In addition, since σ_A is of finite order, we can apply Proposition 2.2 and get that any vertex OISIBO $\tilde{Z}_i$ and any edge OISIBO $\tilde{Z}_j$ between $\tilde{Z}_1$ and $\tilde{Z}_k$ are invariant by σ_A. By Proposition 5.9, for each $\tilde{Z}_i$ and each $\tilde{Z}_j$, Fix $\sigma_A | \tilde{Z}_i$ is a tree and Fix $(\sigma_A | \tilde{Z}_j)$ is (a tree $\times [0, 1]$) .

Note that since the structure map of each attaching map from $\tilde{Z}_{i_0}$ to $\tilde{Z}_{i_1}$ and the restriction of σ_A to $\tilde{Z}_{i_0}$ and $\tilde{Z}_{i_1}$ commute, any point of Fix $(\sigma_A | \tilde{Z}_{i_0})$ is mapped to a point of Fix $(\sigma_A | \tilde{Z}_{i_1})$. Hence Fix σ_A in $\tilde{X}$ is connected. Thus we can find an interval which is fixed by σ_A, and connecting $\tilde{f}(\tilde{x}_\tau)$ and $\tilde{d}$.

We will show that $\pi_2(\tilde{X}) = 0$. Take any continuous map $\varphi : S^2 \to \tilde{X}$. Since $\varphi(S^2)$ is compact, there exists a connected and compact subset P of $\tilde{X}$ which contains a finite number of OISIBO's $\tilde{Z}_i$ of $\tilde{X}$ such that $\varphi(S^2) \subset P$. By Proposition 5.10(ii), for each $\tilde{Z}_i$, $\pi_2(\tilde{Z}_i) = H_2(\tilde{Z}_i) = 0$. Dividing $P = P_1 \cup P_2$ such that P_1 consists of k vertex OISIBO's, $(k-1)$ edge OISIBO's, and $\tilde{Y} \times [0, \frac{1}{2}+\varepsilon]$ and P_2 consists of a vertex OISIBO and $\tilde{Y} \times [\frac{1}{2}-\varepsilon, 1]$, $P_1 \cap P_2 = \tilde{Y} \times [\frac{1}{2}-\varepsilon, \frac{1}{2}+\varepsilon]$ where $\tilde{Y} \times [0,1]$ is an edge OISIBO of $\tilde{X}$, we can show $H_2(P) = 0$ by the induction on k and Mayer-Vietoris exact sequences. Thus $\pi_2(P) = 0$, which gives the fact $[\varphi] = 0$ in $\pi_2(P)$, so is in $\pi_2(\tilde{X})$. Then the triviality of f follows from Proposition 5.7.

The triviality of $O_3(X)$ is derived from showing $\pi_3(\tilde{X}) = 0$ and Proposition 5.8 in the similar mannerr. □

Let M be a 3-orbifold, and X an orbifold composition. We say that two b-continuous maps $f, g : M \to X$ are *C-equivalent* if there is a sequence of b-contininious maps $f = f_0, f_1, \ldots, f_n = g$ from M to X with either f_i is b-homotopic to f_{i-1} or f_i agrees with f_{i-1} on $M - B$ for a certain ballic 3-orbifold $B \subset M$ with $B \cap \partial M$ a discal orbifold or $|B| \cap |\partial M| = \emptyset$. Note that C-equivalent b-continuous maps induce the same homomorphisms $\pi_1(M) \to \pi_1(X)$ up to choices of base points and inner automorphisms.

Remark 5.13. Let f, g be C-equivalent maps from a 3-orbifold M to an orbifold composition X. If $O_i(X)$'s are trivial, $i = 2, 3$, then f and g are b-homotopic.

Lemma 5.14. *Let M be an orbifold, and X an orbifold-composition. Let $p : \tilde{M} \to M$ and $q : \tilde{X} \to X$ be the universal coverings. Suppose dim $\tilde{M} = n$ and $\pi_{n-1}(|\tilde{X}|) = 0$. Let $\tilde{g} : |\tilde{M}| \to |\tilde{X}|$ be a continuous map which satisfies the condition that there exists a homomorphism $\varphi : Aut(\tilde{M}, p) \to Aut(\tilde{X}, q)$ such that for each $\sigma \in Aut(\tilde{M}, p)$, $\tilde{g} \circ \sigma = \varphi(\sigma) \circ \tilde{g}$. Then there exists a continuous map $\tilde{f} : |\tilde{M}| \to |\tilde{X}|$ which satisfies the following:*

1. *There exists a point $\tilde{x} \in |\tilde{M}| - p^{-1}(\Sigma M)$ such that $\tilde{f}(\tilde{x}) \in |\tilde{X}| - q^{-1}(\Sigma X)$.*
2. *There exists an n-ball $B^n \subset |\tilde{M}| - p^{-1}(\Sigma M)$ such that $B^n \cap \sigma(B^n) = \emptyset$ for each $\sigma \in Aut(\tilde{M}, p)$, $\sigma \neq id$, and*

$$\tilde{f}| \left(|\tilde{M}| - \bigcup_{\sigma \in Aut(\tilde{M},p)} \sigma(B^n) \right) = \tilde{g}| \left(|\tilde{M}| - \bigcup_{\sigma \in Aut(\tilde{M},p)} \sigma(B^n) \right). \tag{5.3}$$

3. *For each $\sigma \in Aut(\tilde{M}, p)$, $\tilde{f} \circ \sigma = \varphi(\sigma) \circ \tilde{f}$.*

Proof. The proof is similar to that of [Ta1, Lemma 5.4]. □

Theorem 5.15 (Transversality Theorem)**.** *Let M be a good, compact, connected, orientable 3-orbifold, and X a 3-orbifold composition. Suppose that there exists an edge OISIBO $Y \times [0,1]$ of X, the core Y satisfies that $O_2(X - Y)$ and $O_2(Y)$ are trivial. Then, for any b-continuous map $f : M \to X$, there exists a b-continuous map $g = (|g|, \tilde{g}) : M \to X$ which satisfies the following :*

- **(i)** *g is C-equivalent to f.*
- **(ii)** *Each component of $g^{-1}(Y)$ is a compact, properly embedded, 2-sided, incompressible 2-suborbifold in M.*
- **(iii)** *For properly chosen product neighborhoods $Y \times [-1,1]$ of $Y = Y \times 0$ in X, and $g^{-1}(Y) \times [-1,1]$ of $g^{-1}(Y) = g^{-1}(Y) \times 0$ in M, $|g|$ maps each*

fiber $x \times |[-1,1]|$ *homeomorphically to the fiber* $|g|(x) \times |[-1,1]|$ *for each* $x \in |g^{-1}(Y)|$.

Proof. The proof is similar to that of [Ta1, Theorem 5.5]. $\square$

Corollary 5.16. *In Theorem 5.15, if* $O_i(X)$ *are trivial,* $i = 2, 3$*, then* f *and* g *are b-homotopic by Remark 5.13.*

6. Group representations and splittings of groups

For the contents of the present section we refer to the original paper [C-S].

An isomorphism from a group Π to $\pi_1(G, \mathcal{G})$ is called a *splitting* of Π where $\pi_1(G, \mathcal{G})$ is the fundamental group of a graph of groups $(G, \mathcal{G})$. A splitting is *trivial* if there exists a vertex group which is isomorphic to the whole $\pi_1(G, \mathcal{G})$.

Let Π be a finitely generated group. Take a system of generators $g_1, \ldots, g_n$ for Π. We define a set $R(\Pi) = \{(\rho(g_1), \ldots, \rho(g_n)) \mid \text{a representation } \rho : \Pi \to SL_2(\mathbb{C})\}$. The points of $R(\Pi)$ correspond to the representations of Π in $SL_2(\mathbb{C})$ bijectively, and we often identify such 1-1 corresponding points. For each $g \in \Pi$, we may define a map $\tau_g : R(\Pi) \to \mathbb{C}$ by $\tau_g(\rho) = \mathrm{tr}(\rho(g))$. Let T be the ring generated by all the functions τ_g, $g \in \Pi$. It is finitely generated ([C-S, Proposition 1.4.1]). Thus we can take and fix $\gamma_1, \ldots, \gamma_m \in \Pi$ such that $\tau_{\gamma_1}, \ldots, \tau_{\gamma_m}$ generate T. With those elements we define a map $t : R(\Pi) \to \mathbb{C}^m$ by $t(\rho) = (\tau_{\gamma_1}(\rho), \ldots, \tau_{\gamma_m}(\rho))$, and set $\mathbb{X}(\Pi) = t(R(\Pi))$. For each $g \in \Pi$, there exists a function $I_g : \mathbb{X}(\Pi) \to \mathbb{C}$ which maps $t(\rho)$ to $tr(\rho(g))$. It is a regular function.

Theorem 6.1. ([C-S, Theorem 2.1.1]) *If the group* Π *acts without edge inversions on the tree* T*, then* Π *is isomorphic to* $\pi_1(T/\Pi, \mathcal{G})$ *where* $(T/\Pi, \mathcal{G})$ *is defined in* [C-S, pp.123-124].

Theorem 6.2. ([C-S, Theorem 2.2.1]) *Let* C *be an affine curve contained in* $\mathbb{X}(\Pi)$ *and* $\tilde{C}$ *be a non-singular projective curve uniquely determined by* C*. To each ideal point* $\tilde{x}$ *of* $\tilde{C}$ *one can associate a splitting of* Π *with the property that an element* g *of* Π *is contained in a vertex group if and only if* $\tilde{I}_g$ *does not have a pole at* $\tilde{x}$*. Thus, in particular, the splitting is non-trivial.*

7. Main Theorem

Theorem 7.1. *Let* M *be a good, compact, connected, orientable 3-orbifold without non-separating spherical 2-orbifolds. We assume that the fundamental group of each prime component of* M *is infinite. Suppose that* $\pi_1(M)$ *has a nontrivial finite splitting. Then there exists a non-empty system of essential 2-suborbifolds* $S_1, \ldots, S_n \subset M$ *such that for each component* Q *of* $M - \bigcup_{i=1}^{n} S_i$*,* $\pi_1(Q)$ *is contained in a vertex group.*

Proof. From the hypotheses, $\pi_1(M)$ is isomorphic to $\pi_1(G, \mathcal{G})$, the fundamental group of a graph of groups $(G, \mathcal{G})$. Along the splitting of $\pi_1(M)$, we construct an orbifold composition (of general type) X as follows:

Step a We may assume that M has no spherical boundary components. Take a base point $y_0 \in |M| - \Sigma M$ of M.

Step b By the hypotheses, we can take a prime decomposition of M, each component of which has an infinite fundamental group. Gluing back the prime components by b-continuous maps along the ballic 3-orbifolds attached in capping off, we obtain an OISIBO W with $\pi_1(W) \cong \pi_1(M)$, each particle of which is an irreducible 3-orbifold.

Step c Let G_i, $i \in I$ and H_j, $j \in J$ be the vertex groups and the edge groups of $(G, \mathcal{G})$, respectively. Take the covering OISIBO X_i of W associated with each vertex group G_i of $(G, \mathcal{G})$, and the covering OISIBO Y_j of W associated with each edge group H_j of $(G, \mathcal{G})$. If an edge e_j of G has verteces v_{j_0}, v_{j_1}, then $H_j < G_{j_t}$, $t = 0, 1$. Thus there exist covering maps $p^j_t : Y_j \to X_{j_t}$ which induce monomorphisms $(p^j_t)_* : H_j \to G_{j_t}$, $t = 0, 1$.

Step d The system $X = (X_i, Y_j \times [0,1], p^j_t)_{t=0,1,i \in I, j \in J}$ is a desired orbifold composition with $\pi_1(X) \cong \pi_1(T/\pi_1(M), \mathcal{G}) \cong \pi_1(M)$. Take a base point $x_0 \in |Y_1 \times \frac{1}{2}| - \Sigma(Y_1 \times \frac{1}{2})$ of X.

By Proposition 5.12, $O_i(X)$ are trivial, $i = 1, 2, 3$. By Proposition 5.5, we can construct a b-continuous map $f : M \to X$ which induces an isomorphism $\varphi : \pi_1(M, y_0) \to \pi_1(X, x_0)$.

Note that the set J is finite. For all $j \in J$, take $f^{-1}(Y_j \times \{\frac{1}{2}\})$. By applying Proposition 5.12 for $X - Y_j$, we obtain the fact that $O_2(X - Y_j)$ is trivial. And applying Proposition 5.10 for Y_j, we obtain the fact that $O_2(Y_j)$ is trivial. We have already shown that $O_i(X)$'s are trivial, $i = 2, 3$. With modifications through b-homotopies if necessary, by Theorem 5.15 and Corollary 5.16, each component of $f^{-1}(\bigcup_{j \in J}(Y_j \times \{\frac{1}{2}\}))$ is a compact, properly embedded, 2-sided, incompressible 2-suborbifold in M. If one of such components is boundary parallel, we can reduce the number of components by modifications through b-homotopies. Note that we can make those modifications fixing on some neighborhood of each component of $\bigcup\limits_{j \in J} f^{-1}(Y_j)$, which is already imcompressible. After those modifications, we obtain a system of essential 2-suborbifolds $S_1, \cdots, S_n$ as (not necessarily all) components of $f^{-1}(Y_j \times \{\frac{1}{2}\})$, $j \in J$.

By the construction of $S_1, \cdots, S_n$, for each component Q of $M - \bigcup_{j=1}^n S_j$, $\pi_1(Q)$ is contained in a vertex group of $(G, \mathcal{G})$. □

Corollary 7.2. *Let M be a good, compact, connected, orientable 3-orbifold without non-separating spherical 2-orbifolds. We assume that the fundamental group of each prime component of M is infinite. Suppose that $\pi_1(M)$ acts on a simplicial tree T nontrivially without edge inversions such that $T/\pi_1(M)$ is finite. Then there exists a non-empty system of essential 2-suborbifolds $S_1, \ldots, S_n \subset M$ such that for each component Q of $M - \bigcup\limits_{i=1}^{n} S_i$, $\pi_1(Q)$ fixes a vertex of the tree.*

Proof. By Theorems 6.1 and 7.1. □

Corollary 7.3. *Let M be a good, compact, connected, orientable 3-orbifold without non-separating spherical 2-orbifolds. We assume that the fundamental group of each prime component of M is infinite. Let C be an affine curve contained in $\mathbb{X}(\pi_1(M))$. To each idel point $\tilde{x}$ of $\tilde{C}$ one can associate a splitting of $\pi_1(M)$ with the property that an element g of $\pi_1(M)$ is contained in a vertex group if and only if $\tilde{I}_g$ does not have a pole at $\tilde{x}$. Then there exists a non-empty system of essential 2-suborbifolds*

$S_1, \dots, S_n \subset M$ *such that for each component* Q *of* $M - \bigcup_{i=1}^{n} S_i$, $\pi_1(Q)$ *is contained in a vertex group.*

Proof. By Theorems 6.2 and 7.1. □

REFERENCES

[BMP] M. Boileau, S. Maillot, J. Porti, *Three-Dimensional Orbifolds And Their Geometric Structures*, Panoramas et Synthéses N.15, Société Mathématique de France, (2003).

[C-S] M. Culler and P.B. Shalen, *Varieties of group representations and splittings of 3-manifolds*, Ann. of Math. 117, (1983), pp. 109–146.

[Se] J. P. Serre, *Trees*, Springer-Verlag, Berlin Heidelberg, (1980).

[Sh] P. B. Shalen, *Representations of 3-manifold groups*, in *Handbook of geometric topology*, North-Holland, Amsterdam, 2002, pp. 955-1044.

[Ta1] Y. Takeuchi, *Waldhausen's classification theorem for finitely uniformizable 3-orbifolds*, Trans. of A.M.S. 328, (1991), pp. 151–200.

[T-Y 1] Y. Takeuchi and M. Yokoyama, *PL-least area 2-orbifolds and its applications to 3-orbifolds*, Kyushu J. Math. 55, No. 1, (2001), pp. 19–61.

[T-Y 2] Y. Takeuchi and M. Yokoyama, *The geometric realization of the decompositions of 3-orbifold fundamental groups*, Topology Appl., 95, (1999), pp. 129–153.

[T-Y 3] Y. Takeuchi and M. Yokoyama, *The realization of the decompositions of the 3-orbifold groups along the spherical 2-orbifold groups*, Topology Appl., 124, (2002), pp. 103–127.

[T-Y 4] Y. Takeuchi and M. Yokoyama, *The t-singular homology of orbifolds*, Topology Appl., 153, No. 11, (2006), pp. 1722–1758.

DEPARTMENT OF MATHEMATICS, AICHI UNIVERSITY OF EDUCATION, IGAYA, KARIYA 448-0001, JAPAN
E-mail address: `yotake@auecc.aichi-edu.ac.jp`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, SHIZUOKA UNIVERSITY, OHYA, SURUGA-KU, SHIZUOKA 422-8529, JAPAN
E-mail address: `smmyoko@ipc.shizuoka.ac.jp`